

Defn- A ^(C') curve $\gamma : [a, b] \rightarrow \mathbb{R}^2$ is called closed if $\gamma(a) = \gamma(b)$ and $\gamma'(a) = \gamma'(b)$.

Simple if it is one to one

Convex


Simple closed curve that lie on one side of their tangent lines.

Recall. $t(s) = \langle \cos(\theta(s)), \sin(\theta(s)) \rangle$

$$\theta' = \kappa \quad \left[\theta(s) = \theta_0 + \int_0^s \kappa(t) dt \right]$$

Regard: $t : [a, b] \rightarrow S^1$

$$\theta(b) - \theta(a) = 2k\pi$$



Prop: If $f: [a, b] \rightarrow S^1$, cont. then $\exists$ a
 $f(a) = f(b)$

continuous $\theta: [a, b] \rightarrow \mathbb{R}$ s.t

$$f(t) = \langle \cos(\theta(t)), \sin(\theta(t)) \rangle \quad \forall t \in [a, b]$$

$\theta(t)$ is unique upto adding a multiple of 2π .

i.e

if $\varphi(t)$ is any other polar angle
function then



$$\varphi(t) = \theta(t) + 2k\pi$$

$$\deg(f) = \frac{\theta(b) - \theta(a)}{2\pi}$$

no. of times the domain is wrapped

c.c around the circle

$\deg(\vec{f})$ rotation index of the curve.



Rotation index = 2.

Ex! Do it for a circle with different parametrizations.



$$ds = \frac{1}{K} d\theta \Rightarrow d\theta = K ds$$

$$\int_a^b d\theta = \int_a^b K(s) ds$$

$$\theta(b) - \theta(a) = \text{Total curvature}$$

$$2\pi (\text{rotation index}) = \text{Total curvature}$$

Prop:- Let $f_1, f_2: [a, b] \rightarrow S^1$ be cont. funcs

with $f_1(a) = f_1(b)$ and $f_2(a) = f_2(b)$. If f_1 and f_2 have different degrees that

$$f_1(t_0) = -f_2(t_0) \text{ for some } t_0 \in [a, b].$$

Proof:-

Let $\theta_1, \theta_2: [a, b] \rightarrow \mathbb{R}$ be the angle functions for f_1 & f_2 . Consider

$$\delta(t) = \theta_2(t) - \theta_1(t)$$

Now

$$\begin{aligned} |\delta(b) - \delta(a)| &= |\theta_2(b) - \theta_2(a) - (\theta_1(b) - \theta_1(a))| \\ &\geq 2\pi \end{aligned}$$

∴ degrees are different

Since $\delta(t)$ has a net-change of at least 2π , there must be an odd integer multiple of π between $\delta(a)$ and $\delta(b)$.

Since δ is cont $\Rightarrow$ (IVT) $\exists t_0 \in [a, b]$ s.t


$$\delta(t_0) = (2k+1)\pi, \quad k \in \mathbb{Z}$$

$$\Rightarrow \theta_2(t_0) = \theta_1(b) + (2k+1)\pi$$

$$\Rightarrow f_2(t_0) = -f_1(t_0).$$

Hopf's Theorem

let $\gamma: [a, b] \rightarrow \mathbb{R}^2$ be a simple closed curve. The rotation index of γ is either 1 or -1.



Proof:-

let $\gamma: [a, b] \rightarrow \mathbb{R}^2$ be a simple closed plane curve and let C be

its trace.

let $p \in C$ be a point s.t C is entirely on one side of the tangent line L .

Assume WLOG, γ is parametrized by its arc length and $\gamma(a) = p$.

Consider the triangle


$$T = \{(t_1, t_2) : a \leq t_1 \leq t_2 \leq b\}$$

Define

$\mu: T \rightarrow S'$ as

$$\mu(t_1, t_2) = \begin{cases} \gamma'(t_1) & t_1 = t_2 \end{cases}$$

$$\begin{cases} \frac{\gamma(t_2) - \gamma(t_1)}{\|\gamma(t_2) - \gamma(t_1)\|} & t_1 \neq t_2 \text{ and } (t_1, t_2) \neq (a, b) \\ t - \gamma'(a) & (t_1, t_2) = (a, b). \end{cases}$$



For most inputs $\gamma(t_1, t_2)$ is just a unit vector pointing in the direction from $\gamma(t_1)$ to $\gamma(t_2)$.

Rest ensure γ is continuous.

γ is a loop $\Rightarrow$


$$\lim_{t \rightarrow b-a} \frac{\gamma(t) - \gamma(a)}{b-a-t} = - \lim_{\substack{t \rightarrow b-a \\ t < (b-a)}} \frac{\gamma(b-a) - \gamma(t)}{(b-a)-t}$$

$$\boxed{\gamma(a) = \gamma(a+b-a) = \gamma(b)} = -\gamma'(b-a) = -\gamma'(a)$$

let $\alpha_0: [0,1] \rightarrow T$ be a parametrization of the line segment from (a,a) to (b,b) .

$$\alpha_1: [0,1] \rightarrow T$$

$$(a,a) \rightarrow (a,b) \rightarrow (b,b).$$



$$\alpha_0(t) = (a - at + bt, a - at + bt) \quad \left(\frac{a+b}{2}, \frac{a+b}{2} \right)$$

$$\alpha_1(t) = \begin{cases} (a - 2at + 2at, a - 2at + 2at) \\ (2a - 2at + 2bt - b, b) \end{cases} \quad y - \frac{a+b}{2} = \frac{a+b}{2} - b \quad (x - \frac{a+b}{2} - a)$$

$$(1-2t)(a, a) + 2t(a, b) \\ (2-2t)(a, b) + (2t-1)(b, b)$$

$$y - \frac{a}{2} - \frac{b}{2} = -1 \left(x - \frac{a}{2} - \frac{b}{2} \right)$$

$$y - \frac{a}{2} - \frac{b}{2} = -x + \frac{a}{2} + \frac{b}{2}$$

$$1 - 2(t - \frac{1}{2}) + 2(t - \frac{1}{2})$$

$$y = -x + a + b$$

$$1 - 2t + 1$$

$$\forall s \in [0, 1]$$

$$\alpha_s(t) \stackrel{?}{=}$$

$$(1-t)(a, a) + t(b, b)$$

$$\alpha_s(t) = (a - as + bs, a - as + bs)$$

$$\alpha_1(t) = \{$$

$$\alpha_s(t) = (1-s)\alpha_0(t) + s\alpha_1(t)$$

$(s, t) \mapsto \alpha_s(t)$ is a cont. fn

$$[0,1] \times [0,1] \rightarrow T$$

For each $s \in [0,1]$, let $D(s)$ be the degree of

$$D = \gamma \circ \alpha_s: [0,1] \rightarrow S^1$$


D cont + integer valued $\Rightarrow D$ must be constant

$$D(0) = \gamma \circ \alpha_0$$

$$= \deg(\text{unit tangent of } \gamma)$$

$$= \text{rotation index of } \gamma$$

$$D(0) = D(1) = \gamma \circ \alpha_1$$



(a,a) to (a,b) $\longrightarrow$ one Semicircle (above)

(a,b) to (b,b) $\longrightarrow$ one Semi-circle (below)

$$D(1) = 1 \quad \sigma_2 = -1$$



"Four vertex theorem"

"Every simple closed convex plane curve has at least 4 vertices"


Def:- A point $y(t)$ on the trace is called a vertex if signed curvature is max/min at $y(t)$.

Proof. let C be the trace.

If $K_s = C$ then infinitely many vertices.

Assume vertices of γ are at $p \in C(K_s^{\max})$
and $q \in C(K_s^{\min})$

let L be a line through p and q .



choose arc-length parametrization

$$\gamma: [0, L] \rightarrow \mathbb{R}^2 \text{ s.t. } \gamma(0) = p = \gamma(L)$$

and $\gamma(a) = L$ for some $a \in (0, L)$.

Notice that $K'_s \leq 0$ on $(0, a)$. (max $\rightarrow$ min)

$K'_s \geq 0$ on (a, L) .

WLOG, Assume p is origin and L is x -axis.

Take $y = (x, y)$, y changes sign only at a .

In this case

$K'(t)y(t)$ does not change

Sign on $[0, L]$

$$\Rightarrow \int_0^L y(t) K'(t) dt \neq 0.$$

but

$$y(t)K(t) \Big|_0^L - \int_0^L y'(t)K(t)dt$$

$$\gamma = (x, y) \Rightarrow \gamma' = (x', y')$$

$$\gamma'' = (x'', y'') = K(-y', x')$$

$$- \int_0^L x'' dt$$

$$= -$$